

Supplementary Table 1. General characteristics

Female gender, n (%)	26 (35.6)
Age, year; mean (SD)	66.7 (14.5)
Risk factor, n (%)	
Atrial fibrillation	32 (43.8)
Hypertension	37 (50.7)
Diabetes	17 (23.3)
Hyperlipidemia	19 (26.0)
Current smoking	10 (13.7)
Coronary artery disease	9 (12.3)
Prior stroke	20 (27.4)
Stroke mechanism	
Atherosclerotic	24 (32.9)
Cardioembolic	34 (46.6)
Other	6 (8.2)
Undetermined	9 (12.3)
Initial NIHSS score, mean (SD)	14.2 (5.3)
Occlusion site, n (%)	
M1	43 (58.9)
Distal internal carotid artery	18 (24.7)
Distal internal carotid artery+M1	12 (16.4)
Recanalization therapy, n (%)	
None	9 (12.3)
Intravenous	4 (5.5)
Endovascular	19 (26.0)
Combined	41 (56.2)
Recanalization, n (%)	
Thrombolysis in cerebral infarction (TICI) 0	28 (38.9)
TICI 1	12 (16.7)
TICI 2a	5 (6.9)
TICI 2b	16 (22.2)
TICI 3	11 (15.3)

SD, standard deviation.