

Supplementary Table 2. Statin agent, dose, and time of early statin administration

	Early statins (n=574)	<12 hours (n=191)	12–24 hours (n=187)	24–72 hours (n=196)
Atorvastatin (mg)	541 (94.3)	187 (97.9)	179 (95.7)	175 (89.3)
80	313 (54.5)	139 (72.8)	100 (53.5)	74 (37.8)
40	102 (17.8)	25 (13.1)	42 (22.5)	35 (17.9)
20	32 (5.6)	7 (3.7)	12 (6.4)	13 (6.6)
10	94 (16.4)	16 (8.4)	25 (13.4)	53 (27.0)
Rosuvastatin (mg)	12 (2.1)	1 (0.5)	2 (1.1)	9 (4.6)
20	3 (0.5)	1 (0.5)	1 (0.5)	1 (0.5)
10	9 (1.6)	0 (0.0)	1 (0.5)	8 (4.1)
Simvastatin (mg)	10 (1.7)	2 (1.0)	4 (2.1)	4 (2.0)
20	7 (1.2)	2 (1.0)	3 (1.6)	2 (1.0)
10	3 (0.5)	0 (0.0)	1 (0.5)	2 (1.0)
Pitavastatin (mg)	6 (1.0)	1 (0.5)	1 (0.5)	4 (2.0)
2	6 (1.0)	1 (0.5)	1 (0.5)	4 (2.0)
Pravastatin (mg)	4 (0.7)	0 (0.0)	0 (0.0)	4 (2.0)
40	1 (0.2)	0 (0.0)	0 (0.0)	1 (0.5)
10	3 (0.5)	0 (0.0)	0 (0.0)	3 (1.5)

Values are presented as number (%). High intensity statins were defined as a statin agent and dosage of atorvastatin of 40–80 mg or a dosage of rosuvastatin of 20 mg. Low to moderate intensity statins were defined as all other statin agents/doses.