

**Supplementary Table I.** Probability of a good clinical outcome owing to technically successful angiographic recanalization according to collateral circulation status before intraarterial thrombolysis

Probability of good clinical outcome	Collateral circulation status									
	Good					Poor				
	Estimated	Confidence interval				Estimated	Confidence interval			
		95%	90%	80%	70%		95%	90%	80%	70%
0.8	147.56	-	-	-	32.568	-	-	-	-	-
0.7	225.28	-	57.107	126.801	156.131	-	-	-	-	-
0.6	288.99	159.713	203.168	232.105	246.271	-	-	-	-	-
0.5	347.46	274.959	288.513	301.996	310.497	-	-	-	-	-
0.4	405.93	331.795	341.5	353.124	361.456	30.19	-	-	-	-
0.3	469.64	377.426	387.87	401.212	411.251	93.90	-	-	-	-
0.2	547.36	426.679	439.471	456.268	469.182	171.62	-	-	-	31.942
0.1	664.29	496.487	513.641	536.497	554.274	288.55	9.165	94.807	156.037	187.416

A good clinical outcome was defined as a modified Rankin scale score from 0-2. Collateral circulation was classified as poor (grade 0-1) or good (grades 2-4) according to the American Society of Interventional and Therapeutic Neuroradiology (ASITN) criteria.