

Supplemental Table 3. Clinical outcomes according to the grades of left ventricular diastolic dysfunction

Variables	Grade of left ventricular diastolic dysfunction				P for difference
	Normal	Grade I	Grade II	Grade III	
Outcomes at 3 months after stroke*					
mRS score 2-6	444 (39%)	587 (51%)	162 (49%)	105 (58%)	< 0.01
Recurrent stroke	66 (6%)	81 (7%)	14 (4%)	10 (6%)	0.26
Composite vascular events	70 (6%)	88 (8%)	18 (5%)	11 (6%)	0.37
Outcomes at 1 year after stroke†					
mRS score 2-6	317 (32%)	485 (46%)	127 (41%)	88 (53%)	< 0.01
Stroke recurrence	81 (8%)	93 (9%)	27 (9%)	14 (8%)	0.94
Composite vascular events	89 (9%)	113 (11%)	32 (10%)	16 (9%)	0.55

Composite vascular events include recurrent stroke, myocardial infarction and vascular death after the index stroke.

mRS, modified Rankin Scale.

*Outcomes at 3 months after stroke were based on 2,812 cases for the mRS score and 2,827 cases for vascular events; †Outcomes at 1 year after stroke were based on 2,511 cases for the mRS score and 2,552 cases for vascular events.