

Supplementary Table 3. Detailed results of univariate and multivariate logistic regression models for complete recanalization with modified treatment in cerebral infarction (TICI) 2b or 3 after endovascular revascularization therapy in patients with acute basilar artery occlusion

	Dichotomized analysis			
	Crude OR [CI]	<i>P</i>	Adjusted OR [CI]	<i>P</i>
Age (year)	0.96 [0.90-1.03]	0.256	0.95 [0.88-1.04]	0.293
Male gender	1.44 [0.47-4.42]	0.526	0.49 [0.10-2.39]	0.379
Medical history				
Atrial fibrillation	3.00 [0.84-10.76]	0.092	1.75 [0.29-10.49]	0.538
Hypertension	0.55 [0.16-1.86]	0.338		
Diabetes mellitus	0.90 [0.31-3.09]	0.975		
Hyperlipidemia	0.36 [0.10-1.35]	0.130		
Coronary artery disease	0.64 [0.16-2.61]	0.530		
Previous stroke	0.47 [0.14-1.57]	0.221		
Current smoking	0.47 [0.12-1.82]	0.276		
Stroke informations				
TOAST classification				
Large artery atherosclerosis	Ref.			
Cardioembolism	2.29 [0.70-7.55]	0.173		
Undetermined etiology	2.06 [0.19-22.51]	0.553		
Localization of occlusion				
Caudal to mid-basilar	Ref.			
Distal	1.6 [0.51-5.02]	0.420		
Baseline NIHSS, per 1 point increase	1.00 [0.93-1.07]	0.959	0.95 [0.86-1.05]	0.317
Baseline ASPECT score, per 1 point increase	0.98 [0.75-1.29]	0.906	0.90 [0.64-1.28]	0.563
Clear onset	2.67 [0.81-8.78]	0.107		
Time from stroke onset or LNT to hospital arrival, per 1 hour increase	1.00 [0.996-1.001]	0.336		
Time from stroke onset or LNT to groin puncture, per 1 hour increase	1.00 [0.997-1.001]	0.332	1.00 [0.99-1.00]	0.117
Intravenous tPA, N (%)	1.25 [0.387-4.041]	0.709		
Technical strategies of ERT				
IAF	Ref.			
EMT	10.15 [2.70-38.12]	0.001	23.03 [3.09-171.71]	0.001

TOAST, Trial of ORG 10172 in Acute Stroke Treatment; IQR, interquartile range; LNT, Last Normal Time; NIHSS, National Institutes of Health Stroke Scale; pc-ASPECT, posterior circulation Acute Stroke Prognosis Early Computed Tomography Score; tPA, tissue Plasminogen Activator; IAT, intra-arterial fibrinolysis; EMT, endovascular mechanical thrombectomy