

Supplementary Table 4. Comparisons in favorable outcomes rate according to use of intravenous thrombolysis prior to endovascular treatment in key subgroups using the stabilized Inverse propensity score as weight into logistic regression model

Subgroup	OR (95% CI)	P	P _{het}
Age (yr)			0.820
<80	1.45 (1.11–1.90)	0.007	
≥80	1.55 (0.91–2.64)	0.100	
NIHSS score			0.480
<10	1.56 (0.86–2.83)	0.140	
10–17	1.70 (1.18–2.46)	0.005	
≥18	1.31 (0.89–1.94)	0.170	
ASPECTS			0.041
<5	1.02 (0.47–2.24)	0.950	
5–6	1.92 (0.53–1.56)	0.750	
≥7	1.67 (1.26–2.22)	<0.001	
Site of occlusion			0.580
M2–MCA	1.50 (1.08–2.08)	0.016	
M1–MCA	1.12 (0.58–2.18)	0.730	
Carotid T	1.45 (0.81–2.57)	0.210	
Tandem	1.24 (0.74–2.08)	0.410	
Admission mode			0.500
Mothership	1.47 (0.97–2.24)	0.071	
Drip and ship	1.34 (0.99–1.81)	0.058	
Antiplatelet use			0.160
No	1.25 (0.96–1.63)	0.100	
Yes	1.79 (1.15–2.76)	0.009	

OR, odds ratio; CI, confidence interval; het, heterogeneity; NIHSS, National Institutes of Health Stroke Scale; ASPECTS, Alberta Stroke Program Early CT Score; MCA, middle cerebral artery.