

Supplementary Table 1. Baseline characteristics of stroke patients with intravenous thrombolysis alone or intravenous thrombolysis plus mechanical thrombectomy

Sociodemographic characteristic	IVT alone (n=142)	IVTMT (n=139)	P
Age (yr)	62.8±15.2	63.7±12.7	0.59*
≤70	79 (55.6)	80 (57.6)	0.75 [†]
>70	63 (44.4)	59 (42.4)	
Sex			0.08 [†]
Male	67 (47.2)	80 (57.6)	
Female	75 (52.8)	59 (42.4)	
Comorbidities			
Hypertension	77 (54.2)	64 (47.1)	0.23 [†]
Diabetes mellitus	17 (12.0)	8 (5.9)	0.08 [†]
History of stroke	9 (6.4)	9 (6.8)	0.89 [†]
Hypercholesterolemia	71 (55.0)	58 (48.7)	0.32 [†]
Current or past tobacco use	52 (40.3)	55 (44.7)	0.48 [†]
Coronary disease	20 (14.7)	20 (15.2)	0.92 [†]
Baseline NIHSS score	18 (14–21)	18 (14–20)	0.99 [‡]
ASPECTS at baseline	7 (5–9)	7 (6–8)	0.12 [‡]
Workflow time (min)			
From stroke onset to MR imaging	110 (90–138)	114 (84–138)	0.57 [‡]
From stroke onset to IVT	149 (124–180)	150 (120–175)	0.94 [‡]
From stroke onset to randomization	167 (137–197)	163 (142–192)	0.474 [‡]
Baseline occlusion location (%)			0.047 [§]
ICA	28 (19.7)	15 (10.8)	
M1	114 (80.3)	124 (89.2)	

Values are presented as mean±standard deviation, number (%), or median (interquartile range).

IVT, intravenous thrombolysis; IVTMT, intravenous thrombolysis mechanical thrombectomy; NIHSS, National Institutes of Health Stroke Scale; ASPECTS, Alberta Stroke Program Early CT Score; MR, magnetic resonance; ICA, internal carotid artery.

*Student t-test; [†]Chi-square test; [‡]Median test; [§]Fisher exact test.