

Supplementary Table 14. Net clinical benefit of the combination of death and ICH

Type of diagnosis	All-cause death					ICH				
	Not receiving OACs		Receiving exposure to all OACs		Difference in rate of death (off-on)	Not receiving OACs		Receiving exposure to all OACs		Difference in rate of ICH (on-off)
	PY	Annual rate per 100 PY	PY	Annual rate per 100 PY		PY	Annual rate per 100 PY	PY	Annual rate per 100 PY	
Total cohort	332.8	40.6	239.6	16.3	24.3	315.7	11.4	229.5	5.2	-6.2
Non-traumatic diagnosis	195.2	41.0	113.3	17.7	23.3	185.7	12.4	109.6	3.6	-8.8
Traumatic diagnosis	137.6	40.0	126.3	15.0	25.0	130.0	10.0	119.9	6.7	-3.3
Annual net clinical benefit	Weight=1.5				Weight=1.0	Weight=2.0				
Total cohort	34.1 (33.7–34.5)				30.9 (30.6–31.3)	37.3 (36.9–37.8)				
Non-traumatic diagnosis	37.1 (36.6–37.7)				32.6 (32.2–33.1)	41.7 (41.1–42.3)				
Traumatic diagnosis	30.2 (29.6–30.7)				28.5 (28.0–29.0)	31.9 (31.2–32.5)				

Values are presented as annual net clinical benefit (95% confidence interval).
 ICH, intracranial hemorrhage; OAC, oral anticoagulant; PY, person-year.