Study or subgroup	Mortality Mean	Mortality SD	Mortality Total	No mortality Mean	No mortality SD	No mortality Total	Weight	SMD IV, random, 95% CI
Ahmed et al., 2009	152	22	1,379	151	21	8,096	59.4%	0.05 [-0.01, 0.10]
Molina et al., 2009	176	19	3	153	21	32	4.5%	1.08 [-0.14, 2.29]
Tsivgoulis et al., 2009	162	27	61	156	21	231	36.2%	0.27 [-0.02, 0.55]
Total (95% CI)			**1,443**			**8,359**	**100.0%**	**0.17 [-0.09, 0.44]**

Heterogeneity: Tau^2=0.03; Chi^2=4.94, df=2 (*P*=0.08); I^2=60%
Test for overall effect: Z=1.27 (*P*=0.20)



Study or subgroup	Mortality Mean	Mortality SD	Mortality Total	No mortality Mean	No mortality SD	No mortality Total	Weight	SMD IV, random, 95% CI
Ahmed et al., 2009	83	15	1,375	83	13	8,086	99.8%	0.00 [-0.06, 0.06]
Molina et al., 2009	77	25	3	79	14	32	0.2%	-0.13 [-1.32, 1.05]
Total (95% CI)			**1,378**			**8,118**	**100.0%**	**-0.00 [-0.06, 0.06]**

Heterogeneity: Tau^2=0.00; Chi^2=0.05, df=1 (*P*=0.83); I^2=0%
Test for overall effect: Z=0.01 (*P*=0.99)



Supplementary Figure 11. Forest plots evaluating the association of pre-treatment (A) systolic blood pressure levels and (B) diastolic blood pressure levels with mortality. SMD, standardized mean difference; SD, standard deviation; IV, intravenous; CI, confidence interval.

https://doi.org/10.5853/jos.2018.02369