

Supplementary Table 1. Logistic regression model (dependent variable symptomatic intracranial hemorrhage)

Variable	aOR	95% CI	P
IA UK vs. IA tPA	0.53	0.27–1.04	0.064
Age (per year increase)	1.01	0.97–1.05	0.727
Female sex	0.30	0.095–0.97	0.044*
Admission NIHSS (per point increase)	0.97	0.88–1.06	0.474
Admission ASPECTS (per point increase)	0.78	0.58–1.06	0.110
Symptom-onset/last-seen well to reperfusion (per minute increase)	1.00	0.99–1.00	0.580
TICI (per grade increase)	0.43	0.20–0.94	0.035*
History of diabetes	1.16	0.41–3.24	0.783
History of coronary artery disease	1.25	0.44–3.53	0.672
Anticoagulants			
No anticoagulants	Baseline	Baseline	Baseline
Vitamin K antagonists	1.05	0.11–10.13	0.963
Direct oral anticoagulants	3.08	0.55–17.28	0.201
Antiplatelet medication			
No antiplatelet medication	Baseline	Baseline	Baseline
Aspirin	2.30	0.74–7.09	0.146
Aspirin+other antithrombotic	Did not converge	Did not converge	Did not converge
Admission imaging modality CT vs. MRI	0.52	0.14–1.92	0.328
IV tPA	2.33	0.75–7.17	0.142
Site of intracranial occlusion			
Intracranial ICA	Baseline	Baseline	Baseline
M1	0.27	0.08–0.86	0.026*
M2	0.29	0.07–1.14	0.077

M1/M2, first and second segment of the middle cerebral artery, respectively.

aOR, adjusted odds ratio; CI, confidence interval; IA, intra-arterial; UK, Urokinase; tPA, tissue plasminogen activator; NIHSS, National Institutes of Health Stroke Scale; ASPECTS, Alberta Stroke Program Early CT Score; TICI, thrombolysis in cerebral infarction; CT, computed tomography; MRI, magnetic resonance imaging; IV, intravenous; ICA, internal carotid artery.

*P<0.05.

Supplementary Table 2. Risk of symptomatic intracranial hemorrhage with strata of prior antithrombotic and oral anticoagulant intake

	No sICH	sICH
Antiplatelet medication*		
No antiplatelet medication	199 (91.7)	18 (8.3)
Aspirin	70 (88.6)	9 (11.4)
Aspirin+another antiplatelet medication (e.g., prasugrel)	9 (100)	0 (0)
P		0.59
Oral anticoagulants†		
No oral anticoagulant medication	126 (91.8)	21 (8.2)
Vitamin K antagonists	22 (91.7)	2 (8.3)
Direct oral anticoagulants	21 (84.0)	4 (16.0)
P		0.44

Values are presented as number (%). P-value calculated using Fisher's exact test. sICH, symptomatic intracranial hemorrhage.

*Data available for 308 patients; †Data available for 309 patients.