

Supplementary Table 8. The proportion of patients hospitalized with COVID-19 with concomitant diagnosis of CVT

Variable	No. of centers	CVT with COVID-19	COVID-19 admission	Percentage	95% CI
Overall	154	103	292,080	0.04	0.03–0.05
Asia	40	14	47,253	0.03	0.02–0.05
North America	34	14	102,920	0.01	0.01–0.02
Europe	65	50	110,522	0.05	0.04–0.07
South America	7	12	20,912	0.06	0.03–0.10
Oceania	3	0	68	0	0.00–5.35
Africa	5	13	10,405	0.12	0.07–0.21

COVID-19, coronavirus disease 2019; CVT, cerebral venous thrombosis; CI, confidence interval.

Supplementary Table 9. Mortality as it relates to COVID-19 status during the first year of the COVID-19 pandemic from 120 centers

Variable	Alive	Dead	In-hospital mortality rate (%)	Total
CVT COVID-19 negative	869	41	4.5	910
CVT COVID-19 positive	45	8	15	53
Total CVT admissions	914	49	5.1	963

During the first pandemic year, CVT mortality was higher in patients who were COVID-19 positive compared to COVID-19 negative patients (8/53 [15.0%] vs. 41/910 [4.5%], $P=0.004$).

COVID-19, coronavirus disease 2019; CVT, cerebral venous thrombosis.