


Supplementary Table 8. Medical history and kidney function according to the status of anemia

Anemia on admission	Value (n=10,818)
None (n=8,369)	
Gastroduodenal ulcer	5,687 (68.0)
Hemorrhage of GI tract	448 (5.4)
eGFR (mL/min/1.73 m ²)	
≥90	3,452 (41.2)
60–89	3,659 (43.7)
30–59	1,172 (14.0)
<30	86 (1.0)
Mild (n=1,539)	
Gastroduodenal ulcer	1,157 (75.2)
Hemorrhage of GI tract	107 (7.0)
eGFR (mL/min/1.73 m ²)	
≥90	443 (28.8)
60–89	582 (37.8)
30–59	427 (27.7)
<30	87 (5.7)
Moderate to severe (n=910)	
Gastroduodenal ulcer	718 (78.9)
Hemorrhage of GI tract	117 (12.9)
eGFR (mL/min/1.73 m ²)	
≥90	249 (27.4)
60–89	249 (27.4)
30–59	268 (29.5)
<30	144 (15.8)

GI, gastrointestinal; eGFR, estimated glomerular filtration rate.



Supplementary Figure 1. Study population. CRCS-K, Clinical Research Collaboration for Stroke in Korea; NHIS, National Health Insurance Service.